


A car accelerates from 13 m/s to 42 m/s at a rate of 12 m/s^2 . How long did it take the car to reach the final velocity?



$$a = 12 \text{ m/s}^2$$

$$t = ?$$

$$v_i = 13 \text{ m/s} \quad v_f = 42 \text{ m/s}$$


$$a = \frac{v_f - v_i}{t}$$

$$t = \frac{v_f - v_i}{a}$$

$$= \frac{42 \text{ m/s} - 13 \text{ m/s}}{12 \text{ m/s}^2}$$

$$= 2.42 \text{ s}$$

A car with a mass of 4200 kg and a velocity of 35 m/s slams into a stationary car of mass 2200 kg. If the first car comes to a complete stop after the collision, what is the velocity of the second car?



$$\vec{p} = m\vec{v}$$

$$\vec{p}_{1i} + \vec{p}_{2i} = \vec{p}_{1f} + \vec{p}_{2f}$$

$$m_1 \vec{v}_{1i} + m_2 \vec{v}_{2i} = m_1 \vec{v}_{1f} + m_2 \vec{v}_{2f}$$

$$m_1 v_{1i} = m_2 v_{2f}$$

$$v_{2f} = \frac{m_1 v_{1i}}{m_2}$$

$$= \frac{(4200 \text{ kg})(35 \text{ m/s})}{2200 \text{ kg}}$$

$$= 66.82 \text{ m/s}$$

Find acceleration of second car if collision happens in 0.25 s.


$$a = \frac{v_f - v_i}{t}$$

$$= \frac{66.82 \text{ m/s} - 0 \text{ m/s}}{0.25 \text{ s}}$$

$$= 267.28 \text{ m/s}^2$$

<u>Variable</u>	<u>unit</u>
distance	m
displacement	m
time	s
speed	m/s
velocity	m/s
acceleration	m/s^2 (m/s/s)
momentum	$\text{kg} \cdot \text{m/s}$

A 3500 kg elephant with a velocity of 4 m/s collides with a stationary elephant. The second elephant begins to move at 2.75 m/s. What is the mass of the second elephant?




$$m_1 v_{1i} + m_2 \cancel{v_{2i}} = m_1 \cancel{v_{1f}} + m_2 v_{2f}$$

$$M_1 v_{1i} = M_2 v_{2f}$$

$$M_2 = \frac{m_1 v_{1i}}{v_2}$$

$$= \frac{(3500 \text{ kg})(4 \cancel{\text{m/s}})}{2.75 \cancel{\text{m/s}}}$$

$$= 5090.91 \text{ kg}$$




$$\text{slope} = \frac{\text{rise}}{\text{run}} = \frac{\text{velocity}}{\text{time}} = \text{acceleration}$$

$$= \frac{y_2 - y_1}{x_2 - x_1} = \frac{v_2 - v_1}{t_2 - t_1} = a$$

positive slope \rightarrow velocity increases
so the acceleration is positive

negative slope \rightarrow velocity decreases
so the acceleration is negative

\emptyset slope \rightarrow velocity is constant,
(no) so the acceleration is \emptyset



Line of Best Fit:

- Estimate of data we have
- Goes through points equally spaced above and below
- Get an equation $y = mx + b$

